



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

If $p > n$, then, since there can be but n different ways of voting, n will be the number of different ways the voting may result.

If $p < n$, then since p persons can prepare only p states of the poll, p will be the number of different ways the voting may result.

Also solved by H. C. Whitaker.

GEOMETRY.

Conducted by B.F. FINKEL, Kidder, Missouri. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

40. Proposed by J. C. CORBIN, Pine Bluff, Arkansas.

If R , r , r_1 , r_2 , and r_3 be, respectively, the radii of the circumscribed, inscribed, and escribed circles of a \triangle , prove $r_1 + r_2 + r_3 - r = 4R$.

Solution by M. A. GRUBER. War Department, Washington, D. C.

From any \triangle whose sides are a , b , and c , we obtain $R = \frac{abc}{4\Delta}$,

$$r = \frac{\Delta}{s}, \quad r_1 = \frac{\Delta}{s-a}, \quad r_2 = \frac{\Delta}{s-b}, \quad \text{and} \quad r_3 = \frac{\Delta}{s-c}.$$

$$\begin{aligned} \therefore r_1 + r_2 + r_3 - r &= \frac{\Delta}{s-a} + \frac{\Delta}{s-b} + \frac{\Delta}{s-c} - \frac{\Delta}{s} = \frac{2s^3 - as^2 - bs^2 - cs^2 + abc}{\Delta} \\ &= \frac{s^2[2s - (a+b+c)] + abc}{\Delta} = \frac{abc}{\Delta}. \quad \text{But } \frac{abc}{\Delta} = 4R. \quad \therefore r_1 + r_2 + r_3 - r = 4R. \end{aligned}$$

We might appropriately add a few other combinations of these radii.

$$(1) \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{1}{r}; \quad (2) \quad rr_1r_2r_3 = \Delta^2; \quad (3) \quad Rrr_1r_2r_3 = \frac{abc\Delta}{4}.$$

Solutions of this problem were received from G. I. Hopkins, E. W. Morrell, P. S. Berg, G. B. M. Zerr, F. P. Matz, Cooper D. Schmitt, P. H. Philbrick, J. F. W. Scheffer, John B. Faught, and the Proposer. H. C. Whitaker did not solve the problem but referred to Chauvenet's Geometry and Hallowell's Geometrical Analysis, p. 225.

41. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Find the length (x) of a rectangular parallelopiped $b=5\text{ft.}$, and $h=3\text{ft.}$, which can be diagonally inscribed in a similar parallelopiped $L=83\text{ft.}$, $B=64\text{ft.}$, and $H=50\text{ft.}$

Solution by B. F. BURLERSON, Oneida, Castle, New York, and the PROPOSER.

Let $x = AF$, $y = AE$, $z = P_2D_2$, and $l = O_1P_1$ = the required length of the inscribed rectangular parallelopiped; then, obviously, $x^2 + y^2 = b^2 \dots (1)$, $(L-x)^2 + (B-y)^2$

$$+ (H-z)^2 = l^2 \dots (2),$$

$$x(L-x) = y(B-y) \dots (3),$$

$$\text{and } h\sqrt{[(L-x)^2}$$

$$+ (B-y)^2]} = lz \dots (4).$$

From (3) and (1),

$$4y^4 - 4By^3 + (B^2 - 4b^2 + L^2)y^2$$

$$+ 2Bb^2y = (L^2 - b^2)b^2$$

$$\dots (5); \text{ and this with coefficients numerically ex-}$$

$$\text{pressed, becomes}$$

$$4y^4 - 256y^3 + 10885y^2$$

$$+ 3200y = 171600 \dots (6).$$

Therefore, by *Horner's*

Method of Approximation, we have from (6), $y = 4$; whence $x = 3$. Briefly putting the now known $(L-x)^2 + (B-y)^2 = m^2 = 10000$, we have from (2) and (4), respectively, $m^2 + (H-z)^2 = l^2 \dots (7)$, and $lz = hm \dots (8)$. Therefore, $l^4 - (H^2 + m^2)l^2 + 2Hhml = h^2m^2 \dots (9)$; that is, $l^4 - 12500l^2 + 30000l = 90000 \dots (10)$.

Whence $l = 110.617130324415$ feet.

COR.—Make $H = 0$, and $h = 0$; then the problem becomes: *Find the length of a rectangle of given width inscribed diagonally in a given rectangle.*

After performing obvious operations, we obtain

$$l^4 - (B^2 + 2b^2 + L^2)l^2 + 4BbLl = (B^2 - b^2 + L^2)b^2 \dots (11); \text{ or with the coefficients numerically expressed, we have the equation,}$$

$$l^4 - 11035l^2 + 106240l = 274000 \dots (12).$$

Therefore $l = 100$ feet, which is the length of the diagonally-inscribed rectangle required.

A. H. Bell gets 107.5 feet as a result.

42. Proposed by G. I. HOPKINS, Instructor in Mathematics and Physics in High School, Manchester, New Hampshire.

If the bisectors of two angles of a triangle are equal the triangle is isosceles.

Solution by the PROPOSER.

Let AF and BD bisect the angles of the triangle ABC , and let $AF = BD$.

Draw DE . Make $\angle PDO = \angle PDF$, and $\angle QFN = \angle QFD$.

Draw AH perpendicular to AF and BK perpendicular to BD .

Draw FH through O and DK through N .

